

Hemochromatosis Treatment by Venipuncture Through History

Marija Čuljak¹

¹ Department of Clinical Chemistry, University Hospital Center Sestre Milosrdnice, Zagreb, Croatia

Corresponding author: Marija Čuljak, culjakmarija@gmail.com

Abstract

Hemochromatosis is a hereditary disease caused by the mutation of genes responsible for regulating iron metabolism in the body. The mutation results in increased absorption of iron from food, which is then deposited in various organs and tissues. Due to the excessive decomposition of iron, organs, most commonly liver, heart and pancreas are damaged. The standard therapeutic procedure for the treatment of hereditary hemochromatosis is phlebotomy or venipuncture, which removes excess iron from the blood. The therapeutic procedure is carried out until the iron level returns to the reference interval. Ancient, three thousand years old bloodletting skills are still used to this day. It was considered that the bloodletting establishes a good balance of bodily fluids referred to as eucrasia. Venipuncture was used to treat various diseases, with different amount of blood released, from half a liter to two liters, and sometimes even more. Venipuncture was applied not only for the treatment of existing diseases, but also as a preventive measure. To date, the use of therapeutic venipuncture has remained the gold standard for the treatment of hereditary hemochromatosis.

The aim of this paper is to present a historical review of venipuncture or phlebotomy as a therapeutic procedure for the treatment of hemochromatosis.

(Čuljak M. Hemochromatosis Treatment by Venipuncture Through History. SEEMEDJ 2018; 2(1); 44-48)

Introduction

There are various expressions used to describe the procedure of therapeutic blood sampling, venipuncture, phlebotomy, venesection and bloodletting. A widely used expression across the English-speaking world is "phlebotomy", derived from the Greek word "phleb", which stands for "vein", and "tómos" which means "incision". Phlebotomy is the process of opening a vein by incision in order to take blood for therapeutic treatment or diagnosis. One of the first people who recognized the importance of

veins and arteries was the Chinese Emperor known as the Yellow Emperor or Huang-Di (who ruled China in the 27th century BC). His famous remark was that "everything in the blood is under the control of the heart" (1).

The ancient skill of bloodletting dates back to about three thousand years ago. Ancient Greeks, Egyptians, the peoples of Mesopotamia, the Maya and the Aztecs all knew the practice of blood-shedding, or bloodletting, as a treatment for diseases. This practice began with the Egyptians 1000 years before Christ and was

Received: Jan 3, 2018; revised version accepted: Oct 12, 2018; published: November 27, 2018

KEYWORDS: venipuncture, phlebotomy, hemochromatosis, bloodletting

popular among doctors until the late 19th century. Bloodletting was performed by shavers at the recommendation of a physician. Barbershops that performed the procedure were recognized by a barber's pole with red and white stripes. Through history, bloodletting, especially with the help of leeches, was practiced as a treatment for existing diseases and as a method of prevention. Still, not everything was bad in the treatment by bloodletting. Medicinal leeches were used for improving circulation and blood flow during surgery. In some cultures, these treatments are still used today (2).

According to Hippocrates (460 BC - 380 BC), a person's personality (temperament) is determined by four body fluids (called "humors"): phlegm, yellow bile, black bile and blood. Hippocrates defined health as a good mixture (eucrasia), and disease as a bad mixture of these juices (dyscrasia). He used clinical observations to recognize the symptoms of certain diseases. All four classical elements: fire, earth, water and air were considered present in the blood, and it was believed that the letting of blood would restore the patient's general wellbeing. Blood loss was considered beneficial because it was a way of balancing the fluids. Hippocrates believed that the fluids had to be perfectly balanced in order for the individual to have good health. Any imbalance was treated by bloodletting (3).

In Greece, Galen of Pergamon (129-200 AD), a prominent Greek physician, surgeon and philosopher, first described that not only veins were filled with blood, but arteries as well, which were previously considered to be filled with air. He also believed in the humoral balance. He believed that blood should be released near the diseased part of the body. According to him, fever, termination of organ functions and headache are a result of excess blood, so the surgeon should tie the patient's hands to make the veins swell and become more pronounced in order to release a certain amount of blood (4).

In Hebrew teaching, specifically in the Talmud, it is said that the release of blood should be

carried out on certain days of the week and month. Christianity adopts similar guidelines. Islamic scholars also believe that the release of blood is very useful for the treatment of fevers. Physical signs that required bloodletting were redness, swelling, skin warm to the touch, and any sign of pus. The treatment consisted of making numerous cuts on the body, and letting the blood flow into specially prepared receptacles. However, the treatment would be stopped, if the patient felt weak. Interestingly, there is no data on the number of deaths during these procedures (2,3).

Hemochromatosis

Primary hemochromatosis is a hereditary disease caused by HFE gene mutation on chromosome 6. This is the most common autosomal recessive hereditary disease of the Caucasian population in Northern European countries, with prevalence in Europe reported as 1: 300. It is equally represented amongst both male and female populations. Sickness rarely occurs in the African and African American population, as well as in the Asian population. Men develop a clinical picture more often than women, about 10 times more often, the reason for this being the loss of iron in women during periods, birth and breastfeeding. Hereditary hemochromatosis is divided into four subtypes, and only type 1 is clinically important. A total of 85-90% of patients with confirmed diagnosis of hemochromatosis are homozygotes for mutation C282Y (Y / Y genotype), while about 5% are heterozygotes for mutations C282Y, H63D and / or S65C. In individuals who did not demonstrate mutation C282Y, and who are heterozygotes for S65C and / or H63D, the possibility of developing a clinical picture of inherited hemochromatosis is extremely low (5,6).

Diagnoses of hemochromatosis include laboratory diagnostics, radiological diagnostics and histological treatment. The clinical picture of hemochromatosis includes tiredness, hyperpigmentation (bronze skin color), loss of libido in men, diabetes, joint pain, cirrhosis of the liver and cardiomyopathy.

Laboratory diagnostics show elevated levels of serum iron and ferritin as well as increased transferrin saturation index.

Applying venipuncture in the treatment of hemochromatosis

Hemochromatosis was first described in Paris in 1865 by Armand Trousseau, a French physician, who outlined the disease as a set of symptoms involving diabetes, skin pigmentation, and liver cirrhosis. Trousseau did not name the disease (2).

Davis and Arrowsmith are attributed with the first report on the treatment of hemochromatosis which showed the value of venipuncture. In their report, they state that repeated venipunctures can remove harmful iron. To confirm the diagnosis of hemochromatosis, they carried out a liver biopsy. The first patient was a 69-year-old woman. Between March 1947 and February 1949, 40 liters of blood were drawn from this patient. According to their report, all patients treated by bloodletting showed a general improvement (7).

In 1954, Davey, Foxell and Kamp reported the case of a man at the age of 55 who was hospitalized in October 1950. The patient complained of abdominal pain that had lasted for 2 years, increased tiredness and gray and black pigmentation of the face, neck and upper arm. His liver was enlarged, spleen was not palpable, and free intra-abdominal fluid was found. He had not suffered from previous illnesses except for colon cancer surgery in 1936. He drank plenty of beer daily. The tested urine showed glucose in the urine. Skin biopsy showed iron deposits, especially around the sweat glands. The patient was treated with repeated venesections. Over the course of two years, 31 liters of blood were drawn from the patient. Heavy anemia developed, although the patient showed subjective and objective improvement. The patient was last registered in 1954 when his symptoms were pain in the joints. A report was published, claiming that venesection is an effective hemochromatosis treatment method (8). In 1965, a group of authors (Block, Moore, Wasi and Halby) investigated a

cohort of patients with hemochromatosis. In their research, a liver biopsy was performed prior to venesection, the findings of which suggested that liver lesions arise due to abnormal absorption of iron despite a normal diet (9). In an article published in 1969, Dr. R. Williamson reported that over a period of 5 years, the mortality of patients treated with repeat venesection was 11%, while the mortality of untreated patients was more than 60% (10). Sargent et al. report on two hypotheses regarding hereditary hemochromatosis. One hypothesis is that inherited hemochromatosis results in excessive absorption of iron from food consumed by the patient, which is deposited in the organs, where it causes toxic organ damage. The second hypothesis is that hereditary hemochromatosis is actually a form of cirrhosis that is further complicated by excessive iron from the patient's diet. In eight patients with hereditary hemochromatosis, iron levels were tested before, during and after therapy (venipuncture). In all cases, serum iron levels after therapy were within normal limits, as were other blood indices. This is in favor of the hypothesis that inherited hemochromatosis is a disorder in the mechanism responsible for controlling iron absorption from food (11).

Untreated hemochromatosis can result in death due to cardiac complications. Heart damage is manifested as cardiomyopathy, arrhythmia and heart failure. A period of 12 months is required to restore normal liver function and up to 30 months to fully recover from angina. Continuous and controlled therapy (venipuncture) is needed to restore normal parameters (12).

An early diagnosis of hereditary hemochromatosis has a tremendous significance in preventing the development of diabetes mellitus as well as hepatocellular carcinoma. Niederau et al. claim that patients who received timely therapy prior to the onset of cirrhosis did not develop hepatocellular carcinoma later. Early treatment with venipuncture prevents the formation of cirrhosis and attention should therefore be focused on early diagnosis of hereditary hemochromatosis (13).

A research conducted by a group of authors in Norway shows that the treating of hereditary hemochromatosis by venipuncture leads to a change in the level of some trace elements, including increased absorption of toxic elements such as copper. (14).

Conclusion

Therapeutic venipuncture or phlebotomy is the golden standard for the hemochromatosis therapy. In the procedure, 500 ml of blood containing 200 to 250 mg of iron is extracted. Standard procedures prescribe periodic venipuncture, approximately once a week, with the letting of 400-500 ml of blood, during the first 3 years, followed by 400-500 ml every 1-3 months. Venipuncture is carried out until the levels of iron and ferritin drop to the lower limit of the normal reference interval. Venipuncture can be performed on younger patients 2-3 times a week. The aim is to achieve ferritin values below 50 ng / ml and a ferritin saturation index under 50% in serum. Most patients can tolerate quite well a week of undergoing the procedure, which should be performed until hematocrit falls below 37%. To monitor the therapeutic response, it is recommended to check serum saturation and serum ferritin levels every 2 to 3 months (15).

References

1. History of phlebotomy. Available at: <https://phlebotomycoach.com/resources/history-of-phlebotomy>, Accessed October 25th, 2017.
2. DePalma RG, Hayes VW, Zacharski LR. Bloodletting: past and present. *J Am Coll Surg* 2007;205(1):132-144.
3. Colović N, Leković D, Gotić M. Treatment by bloodletting in the past and present. *Srp Arh Celok Lek* 2016;144:240-248.
4. Long History of Phlebotomy-Bloodletting. Available at: <https://hubpages.com/education/Long-History-of-Phlebotomy>, Accessed October 25th, 2017.
5. Ulvik RJ. Hereditary haemochromatosis through 150 years. *Tidsskr Nor Laegeforen* 2016; 136:2017-2021.
6. Topić E, Primorac D, Janković S. Medicinsko biokemijska dijagnostika u kliničkoj praksi. Medicinska naklada, Zagreb, 2004 (in Croatian).
7. Davis WD Jr, Arrowsmith WR. The effect of repeated bleeding in Hemochromatosis. *J Lab Clin Med.* 1950;36:814-815.
8. Davey DA, Foxell AW, Kemp TA. Treatment of haemochromatosis by repeated venesection. *Br Med J* 1954;2:1511-4.
9. Block M, Moore G, Wasi P, Haiby G. Histogenesis of the hepatic lesion in primary hemochromatosis: with consideration of the pseudo-iron deficient state produced by phlebotomies. *Am J Pathol* 1965;47:89-123.

Treatment of primary hemochromatosis by venipuncture is a safe and effective way of removing blood in order to reduce iron levels. Treatment is a lengthy process, usually lasting several years until iron values return to normal. Venipuncture improves the overall condition of the patient, alleviates the symptoms of fatigue, pain, and skin pigmentation. Regular therapy prevents liver complications, heart disease and diabetes (16).

Despite the advances in technology, bloodletting as a therapeutic procedure has survived to this day as a controlled, supervised, safe and efficient treatment option. In this procedure, it is important to follow all the guidelines for proper blood drawing. The role of medical laboratory personnel in diagnosing and monitoring primary hemochromatosis is vital. Laboratory tests allow the physician to make a proper diagnosis and evaluate the overall medical condition during the course of disease monitoring

Acknowledgement. None.

Disclosure

Funding. No specific funding was received for this study.

Competing interests. None to declare.

10. Crosby WH. Hemochromatosis and venesection. *Br Med J* 1969;4:109-10.
11. Sargent T, Saito H, Winchell HS. Iron absorption in hemochromatosis before and after phlebotomy therapy. *J Nucl Med* 1971;12:660-7.
12. Feely J, Counihan TB. Haemochromatosis presenting as angina and responding to venesection. *Br Med J* 1977;2:681-8.
13. Finlayson ND. Hereditary (primary) haemochromatosis. *BMJ* 1990;301:350-1.
14. Crosby WH. A history of phlebotomy therapy for hemochromatosis. *Am J Med Sci* 1991;301:28-31.
15. Bolann BJ, Distant S, Mørkrid L, Ulvik RJ. Bloodletting therapy in hemochromatosis: Does it affect trace element homeostasis? *J Trace Elem Med Biol.* 2015;31:225-9.
16. Cook LS. Therapeutic phlebotomy: a review of diagnoses and treatment considerations. *J Infus Nurs* 2010;33:81-8.